



A Level Maths The Final Countdown

Week begin- ning	Assignment/ Papers*	Tuesday 1	Tuesday 2	Wednesday	Thursday 1	Thursday 3	Friday
24 th Feb	Assignment 19	Differential Equations	P1	P2			Pure Test 1
2 nd March	Assignment 20	Applied Test 1	Р3	P4	Р5	P6	P7
9 th March	Assignment 21			P8	Pure Test 2	P9	P10
16 th March	OSC 13 Pure 1, Pure 2	Pure Mock on March 16 th	No lessons on Tuesday	M1	M2	M3	M4
23 rd March	OSC 14 Pure 3, Pure α	Pure Test 3	S1	52	53	S4	S5
30 th March	OSC 15 Pure β , Applied 1	Applied Test 2	P1, P2	P3, P4	P5, P6	P7, P8	P9, P10
20 th April	OSC16, OSC17, OSC18, Pure γ , Pure δ , Pure 4, Applied 2, Applied 3	Pure Test 4	S1, S2	S3, S4	S5	M1, M2	M3, M4
27 th April	Pure 5, Pure 6, Pure 7, Applied 4	Pure Test 5	Revision	Revision	Applied Test 3	Revision	Last lesson

The plan for the rest of the course is shown below:

Study leave starts on Monday May 4th. You should carry on completing the papers during Study Leave.

4th May: Pure 8, Pure 9, Pure 10, Applied 5 18th May: Pure 14, Pure 15, Pure 16, Applied 7 1st June: Pure 19, Pure 20, Applied 10 **11th May:** Pure 11, Pure 12, Pure 13, Applied 6 **25th May:** Pure 17, Pure 18, Applied 8, Applied 9 **8th June:** Pure 21, Pure 22, Applied 11, Applied 12

*OSC=Old Specification Core Papers (4 hours) Thursday will continue to be the day to hand in Assignments/Papers.

P1 Proof	S1. Sampling
P2. Algebra And Functions	S2. Data Presentation and Interpretation
P3. Coordinate Geometry	S3. Probability
P4. Sequences And Series	S4. Distributions
P5. Trigonometry	S5. Hypothesis Testing
P6. Exponentials And Logarithms	
P7. Differentiation	M1. Units
P8. Integration	M2. Kinematics
P9. Numerical Methods	M3. Forces
P10. Vectors	M4. Moments

Dates of the exams (please check with your personal timetable) June 3rd: Pure Paper 1 June 10th : Pure Paper 2

June 12th: Applied

How to get 100% in every exam

The key to success is **honest self-assessment** followed by **remedial action**. If you are honest with yourself about how much you understand the work **and** if you take remedial action to improve your weak areas, you will get a grade A* (unless you make lots of expensive errors).

The Mark Scheme Is Your Enemy

When you work through these papers, do <u>NOT</u> use the mark scheme every time you get stuck. Try to work out what to do yourself. You won't have the mark scheme in the real exam! Do the test in 2 hours and then mark it. If you don't get full marks on a question, find out what went wrong (use your Survival Kit or the textbook or any other form of support). Then go back to it a day later and do it again (without looking at the mark scheme). Keep doing this until you can get the question right without help. This is the way you will improve your understanding.

In the mark schemes, the following symbols are used:

M marks: method marks are awarded for 'knowing a method and attempting to apply it'.A marks: Accuracy marks can only be awarded if the relevant method (M) marks have been earned.B marks are for the correct answer (method not necessary)

Support

Over the next three months, please continue to email me if you need support. If you do email me, remember to send a picture of what you have done so far and let me know what it is in the mark scheme that you don't understand.

If you get stuck there are a number of things you can do

- Refer to your Survival Kit
- Watch the videos again
- Use the textbook and read the bits between the exercises
- Try some of the questions in the exercises to get further help
- Drop In With The Doubles
- Look at videos and resources from other websites
 - Exam Solutions: <u>https://www.youtube.com/user/ExamSolutions</u>
 - Mr Hegarty: <u>https://www.youtube.com/user/HEGARTYMATHS/videos</u>
 - Physics and Maths tutor http://www.physicsandmathstutor.com/
 - o MadAsMaths <u>http://www.madasmaths.com/</u>
 - Dr Frost Maths <u>http://www.drfrostmaths.com/</u>
 - Owen's Resource Library <u>https://www.tes.com/teaching-resources/shop/Owen134866</u>

However, the best thing you can do is figure it out yourself.

Formulae To Learn

Make sure you remember **all** the formulae (see later in this booklet) The formulae that you don't need to remember are in the formula book (also see later in this booklet)

How To Study An Exam Paper

DO NOT simply copy out the mark scheme.

Here is why you must *study* the exam papers. By following this programme of exam paper study, you will:

Improve your knowledge of how to solve standard problems. By completing <u>every</u> question from past papers and practice papers you will encounter almost every question that has been put into your real exam.

Improve the accuracy of your algebra... The exam board have told us that the only difference between E grade students and A grade students is that the A grade students make fewer algebraic errors. *Being able to answer the questions is not enough*. You need to be able to answer them <u>without</u> making expensive errors, and this is not something you can learn at the last minute. It takes practice.

Improve the <u>speed</u> of your algebra... Your real exam will be an algebra sprint. It is very important that you get used to the speed required.

Studying an exam paper is not the same as doing an exam paper.

This what you should <u>not</u> do:

- Sit down and complete an exam paper.....
- After an hour you have done all you can do so you mark your work using the mark scheme.
- You realise you've made some errors and you think 'oops! I won't do that in the exam!'
- You read the solutions to the questions you couldn't do and think 'oh I see how to do it now'.
- What has this process done to improve your chance of getting a good grade in the real exam?

Do you have any more knowledge? NO. You have read the solutions to the questions you couldn't do, but this <u>doesn't</u> mean you can actually do them.

Have you improved the accuracy of your algebra? \mathbb{NO}_{\bullet} As soon as you finished the paper you went to the mark scheme. You didn't practice the most important part of an exam – looking for and correcting your error.

Have you improved the speed of your algebra? NO. You didn't try to complete each question in a fixed number of minutes so you still have no idea whether you were going at exam speed.

This is what you should do:

- 1. Complete the exam paper in exam conditions. This means you continue working for two hours and make a real determined effort to find your errors before the time is up.
- 2. Don't use
 - a. The mark scheme
 - b. Your Survival Kit
 - c. Textbooks
 - d. Any other support
- 3. Mark the paper carefully using the mark scheme. Add up the marks and determine the grade.
- 4. Look at all the marks you lost categorising them as being due to
 - a. LU = Lack of understanding (not knowing what to do)
 - b. EE = An expensive error (something that seems silly when you realise what you did)
- 5. Study your mistakes using
 - a. The mark scheme
 - b. Your Survival kit
 - c. Textbooks
 - d. Videos
 - e. etc.
- 6. Wait a day then repeat any question that you lost marks on using the strict timing (number of minutes = number of marks) and looking for your errors **before** you look at the mark scheme.
- 7. Repeat steps 3 and 4 over and over again until you are confident that if any of those questions are in your real exam, you will be able to do them quickly and accurately.

What has this process done to improve your chance of getting an improved grade in the real exam?

Do you have any more knowledge? YES. You kept going back to the harder questions until you could do them, so if these questions come up in your exam you, unlike some other students, can be confident you will know what to do.

Have you improved the accuracy of your algebra? YES. Not only have you practised finding errors during step one of the process, the fact that you have written them down and categorised them will help you to be more aware of the sorts of errors you make and this will help you, unlike other students, to avoid making them in the real exam.

Have you improved the speed of your algebra? YES. Every time you complete a paper (or question) in the correct time you are training yourself to be more comfortable working at the speed needed in the exam. This means that, unlike some other students, you won't have the problem of running out of time.

The key to success is honest self assessment followed by remedial action.

Revision Tips

Multiple attempts and crossing out

"Crossed-out work should be marked unless the candidate has replaced it with an alternative response." "Where a candidate has made multiple responses and indicates which response they wish to submit, examiners should mark this response. If there are several attempts at a question which have not been crossed out, examiners should mark the final answer which is the answer that is the most complete."

Circle or highlight key phrases in questions.

Tick off question parts as you go so you don't leave any parts outKey phrases might include3 d.p.HenceSimplified FractionNearest Integerx is an integer, x>1Express the probability as a %Show – Prove – Verify

Read the key phrases carefully

"Give an exact answer" means leave your answer with a fraction or π or \sqrt{r} or ln or e in it. Don't give a rounded decimal.

"Hence" means use what you have just found out.

"Write down" means that there is no working – this shouldn't take you very long.

"Interpret" means that you need to write a sentence in the context of the question.

Prove or Show

If you're asked to "Prove" or "Show" something, the last line in your working should state the answer. It's not just enough to write "As required" or "Q.E.D." – you must write out the statement.

e.g. $f(x) = 2x^3 - 7x^2 + 4x - 4$. Use the factor theorem to show that (x - 2) is a factor of f(x).

f(2) = 2 x 8 - 7 x 4 + 4 x 4 - 4 = 0∴ (x - 2) is a factor of f(x)Therefore (x-2) is a factor.

Prove, Show, Verify

Prove means LHS $\equiv \cdots \equiv \cdots \equiv RHS$ e.g. prove that $1 + \tan^2 \theta \equiv \sec^2 \theta$ \therefore Proof complete or QED

Don't take any shortcuts with proofs. Write out every step. You may know that $\frac{tan\theta}{sec\theta} = sin\theta$ but if it's part of a proof, you must go through the intermediate step.

Show means use the information to get an answer e.g. Show that x = 1.41 is a solution to the equation $x^2 = 2$ correct to 3 s.f. $\therefore x = 1.41$ is a solution to the equation $x^2 = 2$

Verify means substitute a value

e.g. verify that x = 7 is a solution to $x^3 - 2x - 329 = 0$ When x = 7, $x^3 - 2x - 329 = 7^3 - 2x7 - 329 = 343 - 14 - 329 = 0$ $\therefore x = 7$ is a solution to $x^3 - 2x - 329 = 0$

Scan the paper.

You can do the questions in any order. Start with all the familiar questions first. There may be questions that look unfamiliar. Do them last.

Read The Question

Read the question after you've finished it, to check you've done what it asks you to.

(c) Substitute $x = \frac{1}{10}$ into your binomial expansion from part (a) and hence find an approximate value for $\sqrt{2}$. Give your answer in the form $\frac{p}{q}$, where p and q are integers. (2)

Write on the exam paper

You can write on the exam paper. Draw diagrams to help you (e.g. area under a graph) Do a sketch for those questions that involve tangents and normal. Don't try to hold the information in your head. If in doubt, sketch it out.

Write in black ink only. Don't use tippex or highlighters as your answer paper is scanned.

Attempt every part of every question

If you can't do part (a) don't give up. Make up a value for your answer to part (a) and then use it in subsequent parts of the question to earn method marks.

Beware of taking shortcuts with your working.

It doesn't take that long to write out an intermediate step.

Improve Levels of Accuracy

Always write the full calculator display down first. Then check the level of accuracy required in the question. Assume 3 significant figures if no accuracy stated. Always use <u>unrounded</u> answers in any subsequent calculations. Remember g = 9.8

Avoid basic arithmetic and algebraic errors.

Check your work

In many questions, it's possible to take your answer and substitute values back in to the question.

Use the differentiation and integration buttons with limits.

Know your Calculator!

Angle units

- Pure radians unless you see the degrees sign $^\circ$
- Mechanics normally degrees

Use brackets to tell the calculator the order of operations. Use all the functions

- Statistics (mean, standard deviation)
- Distributions (Normal, Binomial)
- Solving quadratic equations
- Solving simultaneous equations
- Integration & Differentiation

 $\lambda L^2 + 7\lambda L - 4 = 0$

5 - (2 - 2c) = 3 - 2c $3^{2} + 5 = 11$ $2 + 5(-1)^{3} = 7$

Write in the correct space

Write within the space given for each question. Don't do question 5 in question 6's space as the papers are scanned in question by question. If you need additional paper ask for some. Label parts of questions clearly (a), (b), (c) etc.

<u>Timing</u>

Don't spend too long on one question.

Keep a close eye on the time.

Put a watch on your desk rather than keep looking at the clock on the wall. (Buy a cheap watch if you don't have one.)

Make sure you know how many marks for each question and aim for a minute per mark.

Use the correct notation

Integration

$$\int \sin^2 x \, dx = \frac{1}{2}x - \frac{1}{4}\sin 2x + c$$
NOT
$$\int \sin^2 x = \frac{1}{2}x - \frac{1}{4}\sin 2x$$
Differentiation
$$2x^2 - \frac{dy}{dx} = c$$

$$y = 3x^2$$
 $\therefore \frac{dy}{dx} = 6x$ NOT $\frac{d}{dx} = 6x$

<u>Formulae</u>

Learn the formulae.

Quote the general form first then substitute in the correct values. Make it clear what values you are substituting.

e.g.
$$\frac{d}{dx}\left(\frac{f(x)}{g(x)}\right) = \frac{f'(x)g(x) - f(x)g'(x)}{(g(x))^2}$$

 $\frac{d}{dx}\left(\frac{e^{2x}}{\cos x}\right) = \frac{2e^{2x}\cos x - e^{2x}(-\sin x)}{\cos^2 x}$
When $x = 0$, $\frac{dy}{dx} = \frac{2+0}{1} = 2$

Calculator guidance

Calculators may always be used to check answers.

Students should write down any equations that they are solving, so these can be checked and credit given.

We will always include instructions in questions to indicate that the use of a calculator is **not** permitted. Phrases used to signal that calculators should **not** be used include:

- Solutions relying entirely on calculator technology are not acceptable.
- Solutions based entirely on graphical or numerical methods are not acceptable.
- Numerical (calculator) integration/differentiation is not accepted in this question.
- Use algebraic integration/differentiation to ...
- Use algebra to ...
- Show that ...
- Prove that ...

Rate how good you think you are on each of these topics

Торіс	* * * *	***	**	*
P1. Proof				
P2. Algebra And Functions				
P3. Coordinate Geometry				
P4. Sequences And Series				
P5. Trigonometry				
P6. Exponentials And Logarithms				
P7. Differentiation				
P8. Integration				
P9. Numerical Methods				
P10. Vectors				
S1. Sampling				
S2. Data Presentation and Interpretation				
S3. Probability				
S4. Distributions				
S5. Hypothesis Testing				
M1. Units				
M2. Kinematics				
M3. Forces				
M4. Moments				

Learn these formulae (they are in the specification pages 49-52).

Appendix 1: Formulae

Formulae that students are expected to know for A Level Mathematics are given below and will not appear in the booklet *Mathematical Formulae and Statistical Tables*, which will be provided for use with the paper.

Pure Mathematics

Quadratic Equations

 $ax^2 + bx + c = 0$ has roots $\frac{-b \pm \sqrt{b^2 - 4ac}}{2a}$

Laws of Indices

 $a^{x}a^{y} \equiv a^{x+y}$ $a^{x} \neq a^{y} \equiv a^{x-y}$ $(a^{x})^{y} \equiv a^{xy}$

Laws of Logarithms

 $x = a^{n} \Leftrightarrow n = \log_{a} x \text{ for } a > 0 \text{ and } x > 0$ $\log_{a} x + \log_{a} y \equiv \log_{a} (xy)$ $\log_{a} x - \log_{a} y \equiv \log_{a} \left(\frac{x}{y}\right)$ $k \log_{a} x \equiv \log_{a} (x^{k})$

Coordinate Geometry

A straight line graph, gradient *m* passing through (x_1, y_1) has equation $y - y_1 = m(x - x_1)$

Straight lines with gradients m_1 and m_2 are perpendicular when $m_1m_2=-1$

Sequences

General term of an arithmetic progression:

 $u_n = a + (n-1)d$

General term of a geometric progression:

 $u_n = ar^{n-1}$

Page | 10

Mensuration

Circumference and area of circle, radius r and diameter d:

$$C = 2\pi r = \pi d \qquad A = \pi r^2$$

Pythagoras' theorem:

In any right-angled triangle where $a,\,b$ and c are the lengths of the sides and c is the hypotenuse, $c^2\!=\!a^2+b^2$

Area of a trapezium = $\frac{1}{2}(a+b)h$, where a and b are the lengths of the parallel sides and h is their perpendicular separation.

Volume of a prism = area of cross section \times length

For a circle of radius r, where an angle at the centre of θ radians subtends an arc of length s and encloses an associated sector of area A:

$$s = r\theta$$
 $A = \frac{1}{2}r^2\theta$

In the triangle ABC

Sine rule:

$$\frac{a}{\sin A} = \frac{b}{\sin B} = \frac{c}{\sin C}$$
Cosine rule:

$$a^2 = b^2 + c^2 - 2bc \cos A$$
Area
$$= \frac{1}{2}ab \sin C$$

$$\cos^2 A + \sin^2 A \equiv 1$$

$$\sec^2 A \equiv 1 + \tan^2 A$$

$$\csc^2 A \equiv 1 + \cot^2 A$$

$$\sin 2A \equiv 2 \sin A \cos A$$

$$\cos 2A \equiv \cos^2 A - \sin^2 A$$

$$\tan 2A \equiv \frac{2 \tan A}{1 - \tan^2 A}$$

Calculus and Differential Equations

<u>Differentiation</u>		Integration		Vectors
Function	Derivative	Function	Integral	$\left x\mathbf{i} + y\mathbf{j} + z\mathbf{k}\right = \sqrt{\left(x^2 + y^2 + z^2\right)}$
x^n	nx^{n-1}	<i>x</i> ^{<i>n</i>}	$\frac{1}{n+1}x^{n+1}+c, \ n\neq -1$	Statistics
sin kx	$k\cos kx$	cos kx	$\frac{1}{k}\sin kx + c$	The mean of a set of data: $\overline{x} = \frac{\sum x}{n} = \frac{\sum fx}{\sum f}$
cos kx	$-k\sin kx$		h.	The standard Neural variables $Z = X - \mu$ where $X = N(\mu \sigma^2)$
e ^{kx}	ke ^k	sin kx	$-\frac{1}{k}\cos kx + c$	The standard Normal variable: $Z = rac{X-\mu}{\sigma}$ where $X \sim \operatorname{N}ig(\mu,\sigma^2ig)$
		le:	1 2.	Mechanics
$\ln x$	<u>1</u>	e^{kx}	$\frac{1}{k}e^{kx}+c$	Forces and Equilibrium
	x			Weight = mass $\times g$
$f(u) + \sigma(u)$	$f'(x) + \sigma'(x)$	1	1	Friction: $F\leqslant \mu R$
f(x) + g(x)	f'(x) + g'(x)	\overline{x}	$\ln x +c, \ x\neq 0$	Newton's second law in the form: $F = ma$
f(x)g(x)	$\mathbf{f}'(x)\mathbf{g}(x) + \mathbf{f}(x)\mathbf{g}'(x)$	$\mathbf{f}'(x) + \mathbf{g}'(x)$	$\mathbf{f}(x) + \mathbf{g}(x) + c$	Kinematics
f(g(x))	f'(g(x))g'(x)	f'(g(x))g'(x)	f(g(x)) + c	For motion in a straight line with variable acceleration:
*(5(<i>m</i>))	* (B(W))B (W)	Area under a curve $= \int_{a}^{b} y dx \ (y \ge 0)$		$v = \frac{\mathrm{d}r}{\mathrm{d}t}$ $a = \frac{\mathrm{d}v}{\mathrm{d}t} = \frac{\mathrm{d}^2r}{\mathrm{d}t^2}$
		Area under a curve	$a = \int_{a} y \mathrm{d}x \ (y \ge 0)$	$r = \int v \mathrm{d}t \qquad \qquad v = \int a \mathrm{d}t$

Know your way around the formula book (but try to learn these if you can)

Pure Mathematics

Mensuration

Surface area of sphere = $4\pi r^2$

Area of curved surface of cone = πr imes slant height

Arithmetic series

 $S_{-} = \frac{1}{2} n(a+l) = \frac{1}{2} n[2a+(n-1)d]$ Binomial series $(a+b)^{n} = a^{n} + \binom{n}{1} a^{n-1}b + \binom{n}{2} a^{n-2}b^{2} + \dots + \binom{n}{r} a^{n-r}b^{r} + \dots + b^{n} \quad (n \in \mathbb{N})$ where $\binom{n}{r} = {}^{n}C_{r} = \frac{n!}{r!(n-r)!}$

$$(1+x)^n = 1 + nx + \frac{n(n-1)}{1 \times 2} x^2 + \ldots + \frac{n(n-1)\dots(n-r+1)}{1 \times 2 \times \dots \times r} x^r + \ldots \quad (|x| < 1, n \in \mathbb{R})$$

Logarithms and exponentials

 $\log_a x = \frac{\log_b x}{\log_b a}$

 $e^{x \ln a} = a^x$

Geometric series

$$S_n = \frac{a(1-r^n)}{1-r}$$

$$S_{\infty} = rac{a}{1-r}$$
 for $|r| < 1$

Trigonometric identities

$$\sin(A \pm B) = \sin A \cos B \pm \cos A \sin B$$
$$\cos(A \pm B) = \cos A \cos B \mp \sin A \sin B$$
$$\tan(A \pm B) = \frac{\tan A \pm \tan B}{1 \mp \tan A \tan B} \quad (A \pm B \neq (k + \frac{1}{2})\pi)$$
$$\sin A + \sin B = 2 \sin \frac{A + B}{2} \cos \frac{A - B}{2}$$
$$\sin A - \sin B = 2 \cos \frac{A + B}{2} \sin \frac{A - B}{2}$$
$$\cos A + \cos B = 2 \cos \frac{A + B}{2} \cos \frac{A - B}{2}$$
$$\cos A - \cos B = -2 \sin \frac{A + B}{2} \sin \frac{A - B}{2}$$
Diff

Small angle approximations

$$\sin\theta \approx \theta$$
$$\cos\theta \approx 1 - \frac{\theta^2}{2}$$

 $\tan\theta \approx \theta$

where heta is measured in radians

Differentiation

First Principles

$$f'(x) = \lim_{h \to 0} \frac{f(x+h) - f(x)}{h}$$

$\mathbf{f}(\mathbf{x})$	f'(x)
tan kx	$k \sec^2 kx$
sec kx	$k \sec kx \tan kx$
$\cot kx$	$-k \operatorname{cosec}^2 kx$
cosec kx	$-k \operatorname{cosec} kx \operatorname{cot} kx$
$\frac{f(x)}{g(x)}$	$\frac{\mathbf{f}'(x)\mathbf{g}(x) - \mathbf{f}(x)\mathbf{g}'(x)}{\left(\mathbf{g}(x)\right)^2}$
8()	

$$f(x) \qquad \int f(x) dx$$

$$\sec^2 kx \qquad \frac{1}{k} \tan kx$$

$$\tan kx \qquad \frac{1}{k} \ln |\sec kx|$$

$$\cot kx \qquad \frac{1}{k} \ln |\sin kx|$$

$$\csc kx \qquad -\frac{1}{k} \ln |\csc kx + \cot kx|, \quad \frac{1}{k} \ln |\tan(\frac{1}{2}kx)|$$

$$\sec kx \qquad \frac{1}{k} \ln |\sec kx + \tan kx|, \quad \frac{1}{k} \ln |\tan(\frac{1}{2}kx + \frac{1}{4}\pi)|$$

$$f \quad dx \qquad f \quad du$$
Numerical Methods

The trapezium rule:
$$\int_{a}^{b} y \, dx \approx \frac{1}{2} h\{(y_{0} + y_{n}) + 2(y_{1} + y_{2} + \dots + y_{n-1})\}$$
, where $h = \frac{b-a}{n}$

The Newton-Raphson iteration for solving
$$f(x) = 0$$
 : $x_{n+1} = x_n - \frac{f(x_n)}{f'(x_n)}$

Statistics

Probability

 $\mathbf{P}(A') = 1 - \mathbf{P}(A)$

 $P(A \cup B) = P(A) + P(B) - P(A \cap B)$

$$P(A \cap B) = P(A)P(B \mid A)$$

$$P(A \mid B) = \frac{P(B \mid A)P(A)}{P(B \mid A)P(A) + P(B \mid A')P(A')}$$

For independent events A and B,

 $P(B \mid A) = P(B)$ $P(A \mid B) = P(A)$ $P(A \cap B) = P(A) P(B)$

Page | 13

Standard deviation

Standard deviation =
$$\sqrt{(Variance)}$$

Interquartile range =
$$IQR = Q_3 - Q_1$$

For a set of *n* values
$$x_1, x_2, \ldots, x_i, \ldots, x_n$$

$$S_{xx} = \Sigma (x_i - \overline{x})^2 = \Sigma x_i^2 - \frac{(\Sigma x_i)^2}{n}$$

Standard deviation =
$$\sqrt{\frac{S_{xx}}{n}}$$
 or $\sqrt{\frac{\sum x^2}{n} - \overline{x}^2}$ $s = vt - \frac{1}{2}at^2$
 $v^2 = u^2 + 2as$
 $s = \frac{1}{2}(u + v)t$

Discrete distributions

Distribution of X	P(X = x)	Mean	Variance
Binomial $\mathrm{B}(n,p)$	$\binom{n}{x}p^{x}(1-p)^{n-x}$	пр	np(1-p)

Sampling distributions

For a random sample of n observations from $\mathrm{N}(\mu,\,\sigma^2)$

$$\frac{X-\mu}{\sigma / \sqrt{n}} \sim \mathcal{N}(0, 1)$$

_

Statistical tables

The following statistical tables are required for A Level Mathematics: Binomial Cumulative Distribution Function (see page 29) Percentage Points of The Normal Distribution (see page 34) Critical Values for Correlation Coefficients: Product Moment Coefficient (see page 37) Random Numbers (see page 38)

Mechanics

Kinematics

v = u + at

 $s = ut + \frac{1}{2}at^2$

For motion in a straight line with constant acceleration:

Use this page to keep a record of the questions you answer from the Revision Material

Topic	Pack						Notes
	1	2	3	4	5	6	
1 (P1)							
2 (P2)							
3 (P3)							
4 (P4)							
5 (P5)							
6 (P6)							
7 (P7)							
8 (P8)							
9 (P9)							
10 (P10)							
11 (S1)							
12 (S2)							
13 (S3)							
14 (S4)							
15 (S5)							
16 (M1)							
17 (M2)							
18 (M3)							
19 (M4)							

94-A* 87-93 80-86 77-79 Α 73-76 70-72 67-69 В 63-66 60-62 57-59 С 53-56 50-52 47-49 D 43-46 40-42 37-39 Ε 33-36 30-32 20-29 U 10-19 0-9 **OSC 15 OSC 16** <u> 0SC 13</u> **OSC 14 OSC 17 OSC 18** P10 P15 P16 P18 P20 A10 P21 A11 A12 P12 P13 P14 P17 A9 P19 P22 P11 Α7 A8 Ρα Pβ A4 P8 P9 A5 A6 Pγ β A1 P4 A2 A3 P5 P6 P3 Ρ7 Ρ1 P2

A Level Maths Exam Practice Chart

Use this page to keep a record of your tests

Test	Percentage	Grade	Notes
Tracking Test 1			
Tracking Test 2			
Tracking Test 3			
Tracking Test 4			
Tracking Test 5			
Pure Mock			
Statistics Mock			
Mechanics Mock			
Pure Test 1			
Pure Test 2			
Pure Test 3			
Pure Test 4			
Pure Test 5			
Applied Test 1			
Applied Test 2			
Applied Test 3			