

# Mark Scheme (Results) January 2011

GCE

## GCE Mechanics M3 (6679) Paper 1

Edexcel is one of the leading examining and awarding bodies in the UK and throughout the world. We provide a wide range of qualifications including academic, vocational, occupational and specific programmes for employers.

Through a network of UK and overseas offices, Edexcel's centres receive the support they need to help them deliver their education and training programmes to learners.

For further information, please call our GCE line on 0844 576 0025, our GCSE team on 0844 576 0027, or visit our website at [www.edexcel.com](http://www.edexcel.com).

If you have any subject specific questions about the content of this Mark Scheme that require the help of a subject specialist, you may find our **Ask The Expert** email service helpful.

Ask The Expert can be accessed online at the following link:

<http://www.edexcel.com/Aboutus/contact-us/>

January 2011

Publications Code UA026583

All the material in this publication is copyright

© Edexcel Ltd 2011

## General Instructions for Marking

1. The total number of marks for the paper is 75.
2. The Edexcel Mathematics mark schemes use the following types of marks:
  - **M** marks: method marks are awarded for 'knowing a method and attempting to apply it', unless otherwise indicated.
  - **A** marks: Accuracy marks can only be awarded if the relevant method (M) marks have been earned.
  - **B** marks are unconditional accuracy marks (independent of M marks)
  - Marks should not be subdivided.


### 3. Abbreviations

These are some of the traditional marking abbreviations that will appear in the mark schemes.

- bod - benefit of doubt
- ft - follow through
- the symbol  $\checkmark$  will be used for correct ft
- cao - correct answer only
- cso - correct solution only. There must be no errors in this part of the question to obtain this mark
- isw - ignore subsequent working
- awrt - answers which round to
- SC: special case
- oe - or equivalent (and appropriate)
- dep - dependent
- indep - independent
- dp decimal places
- sf significant figures
- \* The answer is printed on the paper
- $\square$  The second mark is dependent on gaining the first mark


January 2011  
Mechanics M3 6679  
Mark Scheme

Question Number	Scheme	Marks
1.	$v \frac{dv}{dx} = 7 - 2x$ $\frac{1}{2}v^2 = 7x - x^2 (+c)$ $x = 0 \quad v = 6 \Rightarrow c = 18$ $v = 0 \quad x^2 - 7x - 18 = 0$ $(x + 2)(x - 9) = 0$ $\therefore x = 9$	M1  M1A1  A1   M1  A1   [6]



Question Number	Scheme	Marks
2. (a)	<p>Mass ratio <math>4m</math>    <math>km</math>    <math>(4+k)m</math></p> <p>Dist from <math>O</math>    <math>\frac{3}{8}r</math>    <math>-\frac{1}{2}r</math>    <math>0</math></p> <p>Moments about <math>O</math>:</p> $\frac{3}{8}r \times 4m = \frac{1}{2}r \times km$ <p><math>k = 3</math></p>	<p>B1 B1</p> <p>M1</p> <p>A1</p> <p>(4)</p>
(b)	 <p><math>\tan 30 = \frac{OG}{r}</math></p> $OG = \frac{\lambda}{(7+\lambda)} \times 2r$ $\frac{1}{\sqrt{3}} = \frac{\lambda}{(7+\lambda)} \times 2r \times \frac{1}{r}$ $7 + \lambda = 2\sqrt{3}\lambda$ $\lambda = \frac{7}{(2\sqrt{3}-1)} \text{ (o.e.) or } 2.84$	<p>B1</p> <p>M1</p> <p>A1</p> <p>A1</p> <p>(4) [8]</p>


Question Number	Scheme	Marks
3.	<p>(a)</p> $\text{Vol} = \pi \int_1^2 y^2 dx = \pi \int_1^2 e^{2x} dx$ $= \frac{1}{2} \pi [e^{2x}]_1^2$ $= \frac{1}{2} \pi [e^4 - e^2]$	<p>M1</p> <p>M1 A1</p> <p>A1</p> <p>(4)</p>
	<p>(b)</p> $\text{C of M} = \frac{\int_1^2 \pi y^2 x dx}{\text{vol}}$ $\int_1^2 e^{2x} x dx = \left[ \frac{1}{2} x e^{2x} \right]_1^2 - \int_1^2 \frac{1}{2} e^{2x} dx$ $= \left[ \frac{1}{2} x e^{2x} \right]_1^2 - \left[ \frac{1}{4} e^{2x} \right]_1^2$ $= \frac{1}{2} \times 2e^4 - \frac{1}{2} \times 1e^2 - \left( \frac{1}{4} e^4 - \frac{1}{4} e^2 \right)$ $= \left( \frac{3}{4} e^4 - \frac{1}{4} e^2 \right)$ $\text{C of M} = \frac{\pi \left( \frac{3}{4} e^4 - \frac{1}{4} e^2 \right)}{\frac{1}{2} \pi (e^4 - e^2)} = 1.656\dots$ <p>= 1.66</p> <p>(3 sf)</p>	<p>M1 A1</p> <p>M1</p> <p>A1</p> <p>M1 A1</p> <p>(6)</p> <p>[10]</p>

Question Number	Scheme	Marks
4.	<p>(a)</p> $x = 5 \sin\left(\frac{\pi t}{3}\right)$ $\dot{x} = 5 \times \frac{\pi}{3} \cos\left(\frac{\pi t}{3}\right)$ $\ddot{x} = -5 \times \left(\frac{\pi}{3}\right)^2 \sin\left(\frac{\pi t}{3}\right)$ $\ddot{x} = -\frac{\pi^2}{9} x \quad (\because \text{S.H.M.})$	<p>M1A1</p> <p>A1</p> <p>(3)</p>
	<p>(b)</p> <p>period = <math>\frac{2\pi}{\frac{\pi}{3}} = 6</math></p> <p>amplitude = 5</p>	<p>B1</p> <p>B1</p> <p>(2)</p>
	<p>(c)</p> <p>... = <math>5 \times \frac{\pi}{3} \cos\left(\frac{\pi t}{3}\right)</math>      or <math> v_{\max}  = a\omega</math></p> <p>max. <math>v = \frac{5\pi}{3}</math></p>	<p>M1</p> <p>A1</p> <p>(2)</p>
	<p>(d)</p> <p>At A <math>x = 2</math>      <math>2 = 5 \sin\left(\frac{\pi t}{3}\right)</math></p> <p><math>\sin \frac{\pi}{3} t = 0.4</math></p> <p><math>t_A = \frac{3}{\pi} \times \sin^{-1} 0.4</math></p> <p>At B <math>x = 3</math>      <math>t_B = \frac{3}{\pi} \times \sin^{-1} 0.6</math></p> <p>time A <math>\rightarrow</math> B = <math>\frac{3}{\pi} \times \sin^{-1} 0.6 - \frac{3}{\pi} \times \sin^{-1} 0.4</math></p> <p>= 0.2215... = 0.22 s    accept awrt 0.22</p>	<p>M1</p> <p>A1</p> <p>A1</p> <p>A1</p> <p>(4)</p> <p>[11]</p>

Question Number	Scheme	Marks
5.	<div style="text-align: center;">  </div> <p>(a)</p> $r = \frac{l}{\sqrt{2}}$ $R(\uparrow) \quad T_a \cos 45 = T_b \cos 45 + mg$ $T_a - T_b = mg \sqrt{2} \quad (1)$ $R(\rightarrow) \quad T_a \cos 45 + T_b \cos 45 = mr\omega^2$ $T_a \times \frac{1}{\sqrt{2}} + T_b \times \frac{1}{\sqrt{2}} = m \frac{l}{\sqrt{2}} \omega^2$ $T_a + T_b = ml\omega^2 \quad (2)$ $T_a - T_b = mg \sqrt{2} \quad (1)$ $2T_a = m(l\omega^2 + g\sqrt{2})$ $T_a = \frac{1}{2}m(l\omega^2 + g\sqrt{2})$ $T_b = ml\omega^2 - T_a$ $= \frac{1}{2}m(l\omega^2 - g\sqrt{2})$	<p>B1</p> <p>M1A1</p> <p>M1A1</p> <p>M1</p> <p>A1</p> <p>A1</p> <p>(8)</p>
(b)	$T_b > 0 \quad \frac{1}{2}m(l\omega^2 - g\sqrt{2}) > 0$ $\omega^2 > \frac{g\sqrt{2}}{l} \quad *$	<p>M1</p> <p>A1</p> <p>(2)</p> <p>[10]</p>



Question Number	Scheme	Marks
6. (a)	<div style="text-align: center;">  </div> <p>length <math>AP = \text{length } BP = \frac{5}{4}l</math></p> $T_a = T_b = \frac{kmg \left(\frac{1}{4}l\right)}{l} = \frac{1}{4}kmg \quad (\text{or } T = \dots)$ $R(\uparrow) \quad T_a \cos \theta + T_b \cos \theta = 3mg \quad (\text{or } 2T \cos \theta = 3mg)$ $\frac{1}{4}kmg \times \frac{3}{5} + \frac{1}{4}kmg \times \frac{3}{5} = 3mg \quad \left( \text{or } \frac{1}{2}kmg \times \frac{3}{5} = 3mg \right)$ $\frac{3}{10}kmg = 3mg$ $k = 10 \quad *$	<p>B1</p> <p>M1A1</p> <p>M1A1</p> <p>A1</p> <p>A1</p> <p style="text-align: right;">(7)</p>
(b)	<div style="text-align: center;">  </div> <p>initial extn <math>= \frac{13}{5}l - l = \frac{8}{5}l</math></p> $\text{E.P.E. lost} = 2 \times \frac{\lambda x^2}{2l} = 2 \times \frac{10mg}{2l} \left(\frac{8l}{5}\right)^2 = \frac{128mgl}{5}$ $\text{P.E. gained} = 3mg \times \frac{12l}{5} = \frac{36mgl}{5}$ $\frac{1}{2} \times 3mv^2 + \frac{36mgl}{5} = \frac{128mgl}{5}$ $v^2 = \frac{256gl}{15} - \frac{72gl}{15}$ $v = \sqrt{\left(\frac{184}{15}gl\right)}$	<p>B1</p> <p>M1A1</p> <p>M1A1</p> <p>A1</p> <p style="text-align: right;">(6) [13]</p>

Question Number	Scheme	Marks
7.	<div style="text-align: center;">  </div> <p>(a)</p> $mgl(\cos \alpha - \cos \theta) = \frac{1}{2}mv^2 - \frac{1}{2}mu^2$ $v^2 = u^2 + 2gl(\cos \alpha - \cos \theta) \quad *$	<p>M1A1=A1</p> <p>A1</p> <p>(4)</p>
(b)	$\cos \alpha = \frac{3}{5} \quad v^2 = 2gl\left(\frac{3}{5} - \cos \theta\right) + u^2$ <p>At top <math>\theta = 360^\circ \quad v^2 = 2gl\left(\frac{3}{5} - 1\right) + u^2</math></p> $v^2 > 0 \quad -2gl \times \frac{2}{5} + u^2 > 0$ $u^2 > \frac{4gl}{5}$ $u > 2\sqrt{\frac{gl}{5}} \quad *$	<p>M1A1</p> <p>M1</p> <p>A1</p> <p>(4)</p>

Question Number	Scheme	Marks
(c)	<p>Equation of motion along radius at lowest point:</p> $T_1 - mg = \frac{mv^2}{l}$ $\theta = 180 \quad v^2 = 2gl\left(\frac{3}{5} + 1\right) + u^2$ $v^2 = \frac{16gl}{5} + u^2$ $T_1 = \frac{m}{l}\left(\frac{16gl}{5} + u^2\right) + mg$ $= \frac{21mg}{5} + \frac{mu^2}{l}$ <p>At highest point:</p> $T_2 + mg = \frac{mv^2}{l}$ $\theta = 360 \quad T_2 = 2mg\left(-\frac{2}{5}\right) + \frac{mu^2}{l} - mg$ $T_2 = \frac{mu^2}{l} - \frac{9mg}{5}$ $T_1 = 5T_2$ $\frac{21mg}{5} + \frac{mu^2}{l} = 5\left(\frac{mu^2}{l} - \frac{9mg}{5}\right)$ $\frac{66mg}{5} = \frac{4mu^2}{l}$ $u^2 = \frac{33gl}{10} \quad *$	<p>M1A1</p> <p>M1</p> <p>A1</p> <p>M1</p> <p>M1</p> <p>A1</p> <p>M1</p> <p>A1</p> <p>(9) [17]</p>

Further copies of this publication are available from  
Edexcel Publications, Adamsway, Mansfield, Notts, NG18 4FN

Telephone 01623 467467  
Fax 01623 450481

Email [publications@linneydirect.com](mailto:publications@linneydirect.com)

Order Code UA026583 January 2011

For more information on Edexcel qualifications, please visit [www.edexcel.com/quals](http://www.edexcel.com/quals)

Edexcel Limited. Registered in England and Wales no.4496750  
Registered Office: One90 High Holborn, London, WC1V 7BH